

Liposomal vs Conventional Oral Iron After Sleeve Gastrectomy

NCT07366502

Status	RECRUITING
Phase	Not Applicable
Sponsor	Medeni ^ermet
Enrollment	70 participants

Key Eligibility Criteria

Inclusion (7)

- Age between 18 and 65 years
 - History of laparoscopic sleeve gastrectomy performed at least 6 months prior to enrollment
 - Diagnosis of iron deficiency anemia, defined as:
 - Hemoglobin ≤ 12 g/dL in women or ≤ 13 g/dL in men, and
 - Serum ferritin ≤ 30 ng/mL or transferrin saturation (TSAT) $\leq 20\%$
- ... and 2 more (see full listing online)

Exclusion (10)

- Active gastrointestinal disease (e.g., inflammatory bowel disease, peptic ulcer disease, or gastrointestinal bleeding)
 - Chronic kidney disease with estimated GFR ≤ 30 mL/min/1.73 m²
 - Use of oral or parenteral iron, vitamin B12, or folate supplementation within the previous 3 months
 - Iron overload or hemochromatosis (serum ferritin > 300 ng/mL)
 - Known allergy or intolerance to iron preparations
- ... and 5 more (see full listing online)

Locations (1 total)

Goztepe Prof.Dr. Suleyyman Yalçın City Hospital, Istanbul, kadıköy, Turkey (Türkiye)