

Comparison of the Efficacy of Perineal and Intravaginal Electrical Stimulation in Women With Idiopathic Overactive Bladder

NCT07366918

Status	RECRUITING
Phase	Not Applicable
Sponsor	Pamukkale University
Enrollment	63 participants

Key Eligibility Criteria

Inclusion (5)

- Women over the age of 18 with a clinical diagnosis of idiopathic OAB
- Intolerant or unresponsive to antimuscarinic or oral β_3 adrenoceptor agonist (mirabegron) medications and not using them for at least 4 weeks
- Ability to understand the procedures, advantages, and potential side effects
- Ability to provide written, informed consent
- Pelvic floor muscle (PFM) strength of 3/5 or higher (modified Oxford scale, minimum: 0 - maximum: 5)

Exclusion (11)

- Women with pure stress urinary incontinence
- History of conservative treatment for OAB within the last 6 months (MI, IVES, perineal ES, etc.)
- Pregnant or planning to become pregnant at the time of the study
- Diagnosis of vaginal infection, urinary tract infection, or cancer
- Women with urinary incontinence within the last 3 months Those who have undergone urogynecological surgery
- ... and 6 more (see full listing online)

Locations (1 total)

Pamukkale University, Denizli, Turkey (Türkiye)