

The Analgesic Efficacy and Safety of Loxoprofen Sodium Patch in Patients With Herpes Zoster

NCT07378059

Status	RECRUITING
Phase	Not Applicable
Sponsor	Beijing Tiantan Hospital
Enrollment	750 participants

Key Eligibility Criteria

Inclusion (6)

- \. Ages more than 18 years;
- \. Patients with onset of HZ rash less than 90 days;
- \. Experiencing moderate to severe HZ pain with an average pain score of at least 4 on a Numeric Rating Scale (NRS, 0 = no pain, 10 = worst possible pain);
- \. Aspartate aminotransferase and alanine aminotransferase levels less than twice the upper limit of normal;
- \. Estimated glomerular filtration rate of 30 mL/min per 1.73 m² or higher;
- ... and 1 more (see full listing online)

Exclusion (5)

- \. History of taking loxoprofen sodium patch;
- \. Patients with evidence of cutaneous or visceral dissemination of HZ infection (cutaneous dissemination is defined as more than 20 discrete lesions outside adjacent dermatomes) or ocular involvement of HZ;
- \. History of intolerance or hypersensitivity to any active components or excipient of the loxoprofen sodium patch;
- \. History of systemic immune diseases, organ transplantation, or cancers;
- \. Pregnancy or breastfeeding.

Locations (1 total)

Beijing Tiantan Hospital, Beijing, Beijing 100070, Beijing, China