

# The Study is Being Conducted to Find a New Treatment for Patients Suffering From Interstitial Cystitis Without Hunner's Lesions.

NCT07391306

---

Status	RECRUITING
Phase	Phase 2, Phase 3
Sponsor	JSC NextGen
Enrollment	180 participants

## Key Eligibility Criteria

---

### Inclusion (7)

- Written informed consent prior to any study-related procedures.
  - Women aged 18 to 65 years, inclusive.
  - A confirmed diagnosis of painful bladder syndrome/interstitial cystitis, characterized by pelvic pain associated with filling or emptying the bladder, frequent urination day and night, with symptoms lasting at least 12 months, with the exclusion of a urinary tract infection or other disease that could cause such symptoms.
  - Absence of Hunner's lesions of the bladder during cystoscopy with hydrobougenage of the bladder (2 minutes, 80 cm H2O).
  - A glycated hemoglobin level within the normal range for the patient's age group.
- ... and 2 more (see full listing online)

### Exclusion (15)

- Any condition that prevents intravesical administration of medications.
  - Received medication or non-medication therapy, including physical therapy, for BPS/IC within 1 month prior to study inclusion.
  - Instillation of any pharmacological agent into the bladder less than 1 month prior to screening.
  - History of chronic drug or alcohol abuse.
  - Pregnant women, planning to become pregnant during the study, or currently breastfeeding.
- ... and 10 more (see full listing online)

## Locations (1 total)

---

Botkin Hospital, Moscow, Russia