

Efficacy and Safety of ZKARE® Gel and Its Accessory ZKAPI® in the Treatment of Chronic Ulcers.

NCT07406152

Status	RECRUITING
Phase	Not Applicable
Sponsor	i+Med S.Coop.
Enrollment	134 participants

Key Eligibility Criteria

Inclusion (11)

- Adult patients (≥18 years).
 - Presence of at least one chronic hard-to-heal ulcer, including any of the following etiologies: Grade II, III or IV pressure ulcers; neuropathic diabetic foot ulcers (grade I, II or III), Wagner Grade I-III or Texas 0, I, II, III / Stage B; class C6/C6r venous leg ulcers measured with CEAP scale ; neuro-ischaemic ulcers with adequate distal perfusion (ABI/ITB ≤ 0.7-0.89; mild PAD); mixed-etiology venous ulcers and any chronic lower-limb lesion with palpable distal pulses.
 - Ulcer size between 1 cm² and 100 cm².
 - Ability and willingness to comply with all study visits and procedures according to the clinical protocol.
 - Ability to understand and sign the informed consent, obtained before performing any protocol-specific procedures.
- ... and 6 more (see full listing online)

Locations (1 total)

Araba University Hospital, Vitoria-Gasteiz, Álava, Spain