

Optimal Time for Reperfusion in Acute Pulmonary Embolism

NCT07436702

Status RECRUITING
Sponsor Azienda ULSS 8 Berica
Enrollment 500 participants

Key Eligibility Criteria

Inclusion (4)

- Age ≥ 18 years
- Patients with a diagnosis of intermediate-high or high pulmonary embolism confirmed by computed tomography pulmonary angiography (CTPA), pulmonary angiography, or echocardiography (evidence of a floating thrombus in the right atrium or right ventricle, or proximal visualization of thrombus in the pulmonary artery).
- Classified as high-risk or intermediate-high-risk according to the 2019 European Society of Cardiology guidelines.
- Provision of informed consent for the processing of personal data.

Exclusion (3)

- Age < 18 years
- Patients in whom a diagnosis of acute high-risk or intermediate-high-risk pulmonary embolism has been excluded.
- Inability to provide informed consent.

Locations (7 total)

UO Unità di Cure Intensive Cardiologiche e Cardiologia 1, Ospedale Niguarda, Milano, Milan, Milano, Italy
UOC di Cardiologia, Ospedali Riuniti Padova Sud, Monselice, Italia, Monselice, Padova, Italy
IRCCS Policlinico San Matteo, Pavia, Pavia, Italy
... and 4 more locations