

A Large Language Model-based Chatbot for Alcohol Reduction in Patients With Metabolic Alcohol-Related Liver Disease

NCT07450144

Status	RECRUITING
Phase	Not Applicable
Sponsor	The University of Hong Kong
Enrollment	50 participants

Key Eligibility Criteria

Inclusion (6)

- Hong Kong residents aged 18 years or above
- Diagnosed of MASLD:
- Evidence of hepatic steatosis (fat accumulation in the liver) confirmed by imaging techniques such as ultrasound, computed tomography (CT), magnetic resonance imaging (MRI), or liver biopsy
- Presence of metabolic dysfunction, indicated by at least one of the following:
- Overweight or obesity (body mass index ≥ 25 kg/m²),
- Type 2 diabetes mellitus, Dyslipidemia (elevated triglycerides ≥ 150 mg/dL) or low high-density lipoprotein (HDL) cholesterol (< 40 mg/dL for men, < 50 mg/dL for women), Hypertension (blood pressure $\geq 130/85$ mmHg or use of antihypertensive medication).

... and 1 more (see full listing online)

Exclusion (2)

- Diagnosed with mental disease or cognitive impairments, or
- Participating in other ongoing clinic trials

Locations (1 total)

Hong Kong Queen Mary Hospital, Hong Kong, Hong Kong