

# Left Atrial Appendage Closure With Versus Without Pulsed Field Ablation in Atrial Fibrillation Patients With Mild Symptoms and High Stroke Risk

NCT07453940

---

<b>Status</b>	RECRUITING
<b>Phase</b>	Not Applicable
<b>Sponsor</b>	Sir Run Run Shaw Hospital
<b>Enrollment</b>	50 participants

## Key Eligibility Criteria

---

### Inclusion (5)

- Age ≥ 18 years old.
- Subjects diagnosed with persistent AF with duration more than 3 months.
- Subjects with AFEQT score >70 .
- Subjects with CHA2DS2-VA score ≥2.
- Subjects who are willing and capable of providing ICF and participating in all testing associated with this study.

### Exclusion (15)

- AF that is secondary to electrolyte imbalance, thyroid disease, alcohol, or other reversible/non-cardiac causes.
- Subjects with the history of AF ablation, LAA surgically closed or otherwise excluded or the LAA anatomy does not accommodate a Closure Device.
- Left atrial anteroposterior diameter ≥ 5.5 cm.
- Heart failure with a NYHA III/IV and/or LVEF <35% within 3 months prior to the procedure.
- Any of the following events within 90 days of the Consent Date:  
... and 10 more (see full listing online)

## Locations (1 total)

---

Sir Run Run Shaw Hospital, Hangzhou, Zhejiang, China