

# The Efficacy and Safety of Liposomal Bupivacaine for Transversus Abdominis Plane Block in Relieving Postoperative Pain After Laparoscopic Surgery in Gynecologic Oncology

NCT07458295

---

<b>Status</b>	RECRUITING
<b>Phase</b>	Not Applicable
<b>Sponsor</b>	Beijing Tiantan Hospital
<b>Enrollment</b>	110 participants

## Key Eligibility Criteria

---

### Inclusion (5)

- Patients scheduled for elective Laparoscopic Surgery in Gynecologic Oncology under general anesthesia;
- Ages 18 to 64 years old
- American Society of Anesthesiologists (ASA) physical status of I-III
- Glasgow Coma Scale (GCS) score of 15
- Patients must be able to understand the nature and potential personal consequences of the clinical trial, signing of the informed consent form.

### Exclusion (14)

- History of chronic pain syndrome of any cause.
- Patients with heart conduction block (sinus block or atrioventricular block).
- Patients with unstable coronary artery disease.
- Patients with gastric ulcer or gastric bleeding.
- Patients with diabetes and are being treated with insulin.
- ... and 9 more (see full listing online)

## Locations (1 total)

---

Beijing Tiantan Hospital, Beijing, Beijing Municipality, China