

Transcranial Doppler for Internal Carotid Artery Stenting (TCD-ICAS)

NCT07462546

Status	RECRUITING
Phase	Not Applicable
Sponsor	General Hospital of Shenyang Military Region
Enrollment	232 participants

Key Eligibility Criteria

Inclusion (6)

- Age ≥ 40 years old;
- Patients with severe stenosis of the internal carotid artery;
- Planned for carotid artery stent;
- Modified Rankin Scale score ≤ 2;
- The subject has adequate temporal bone windows to undergo TCD, and the blood flow signal of the middle cerebral artery is detectable.;

... and 1 more (see full listing online)

Exclusion (6)

- Comorbid conditions with severe infections or severe diseases of the liver, kidney, hematopoietic system, endocrine system, etc.;
- Occurrence of intracranial hemorrhage (cerebral parenchymal hemorrhage, subarachnoid hemorrhage, subdural/epidural hemorrhage) within 90 days prior to enrollment;
- Severe hematological disorders or significant coagulation abnormalities;
- Pregnant or breastfeeding women;
- Participation in another clinical trial within the past 3 months or ongoing participation;

... and 1 more (see full listing online)

Locations (1 total)

Department of Neurology, General Hospital of Northern Theater Command, Shenyang, China