

# Preoperative Clindamycin Vaginal Cream Versus Vaginal Cleansing With Povidone-iodine on Postoperative Infections After Cesarean Section

NCT07474194

---

<b>Status</b>	RECRUITING
<b>Phase</b>	Phase 4
<b>Sponsor</b>	Beni-Suef University
<b>Enrollment</b>	150 participants

## Key Eligibility Criteria

---

### Inclusion (4)

- female participants with singleton pregnancies scheduled for elective cesarean sections at term, specifically between 37 and 40 weeks of gestation
- free from any medical disorders.
- The participants' ages ranged from 18 to 40 years
- body mass index (BMI) ranged from 18 to 30 kg/m<sup>2</sup>

### Exclusion (7)

- Cases of urgent or emergent CS.
- Abnormal vaginal secretion (bad smelling yellowish secretion, whitish cheesy discharge with pruritus, bloody discharge; etc...).
- Women with medical disorders such as pre-gestational diabetes, anemia, hypertension, preeclampsia, or immunosuppressant pregnant women.
- Women who had a skin infection adjacent to the operative site.
- Allergy to topical povidone-iodine or clindamycin
- ... and 2 more (see full listing online)

## Locations (1 total)

---

Beni suef university, Ban+ Suwayf, Egypt