

Efficacy of Chemically Distinct Nonsteroidal Anti-Inflammatory Drugs (NSAIDs) and Pain Phenotypes in Adhesive Capsulitis

NCT07493226

Status	RECRUITING
Phase	Not Applicable
Sponsor	Konya Beyhekim Training and Research Hospital
Enrollment	120 participants

Key Eligibility Criteria

Inclusion (4)

- Being between 18-75 years of age
- Having shoulder pain that has lasted for at least one month and being diagnosed with primary frozen shoulder
- Having a VAS pain level e 4/10
- Having e30 degrees of range of motion restriction in at least two planes (flexion, abduction, or external rotation)

Exclusion (11)

- Patients with a history of shoulder trauma, those with musculoskeletal disorders characterized by pain and loss of function in the affected extremity (such as lateral epicondylitis, flexor tendon injury, de Quervein's tenosynovitis)
- Presence of neurological involvement such as stroke, brachial plexus injury, Parkinson's disease, and cervical spine injury with or without radiculopathy
- History of shoulder surgery, malignancy or tumor in the shoulder, those who have undergone shoulder manipulation
- Presence of shoulder arthritis, rotator cuff tear or other shoulder injuries, thoracic outlet syndrome
- Systemic disease affecting the shoulder region, severe degeneration or trauma (e.g., osteoarthritis, rheumatoid arthritis, history of labrum or articular cartilage injuries), inflammatory rheumatic diseases
- ... and 6 more (see full listing online)

Locations (1 total)

Konya Beyhekim Training and Research Hospital, Konya, Selçuklu, Turkey (Türkiye)