

Preserflo MicroShunt Versus Trabeculectomy

NCT07496957

Status	RECRUITING
Phase	Not Applicable
Sponsor	Medical University of Bialystok
Enrollment	100 participants

Key Eligibility Criteria

Inclusion (6)

- >21 years of age
 - An established diagnosis of: Primary open angle glaucoma, Normal tension glaucoma, Pigment dispersion glaucoma (PDG) or Pseudoexfoliative glaucoma (PEXG)
 - Inadequately controlled on maximum tolerated medical therapy
 - Mean Deviation (MD) ≤ -3
 - Intraocular pressure of 14-28 mmHg
- ... and 1 more (see full listing online)

Exclusion (11)

- An established diagnosis of: Closed-angle glaucoma or Secondary open-angle glaucoma (besides PDG and PEX)
 - Lens status: Aphakic patients or Anterior chamber intraocular lens
 - Previous procedures: Glaucoma shunt/valve/ surgery or cyclodestructive procedure, Selective laser trabeculoplasty within the past 3 months, Incisional ophthalmic surgery involving the conjunctiva within the past 3 months, Clear corneal cataract or trabecular meshwork surgery conducted within the past 6 months
 - Presence of intraocular silicone oil
 - No light perception vision
- ... and 6 more (see full listing online)

Locations (1 total)

Medical University of Bialystok, Bialystok, Polska, Poland