

Comparison of the Effects of Oral Hygiene Regimens on Clinical, Immunomodulatory, and Microbial Outcomes and Oral Tolerance in People With Gingivitis

NCT07501455

| | |
|-------------------|-------------------|
| Status | RECRUITING |
| Phase | Not Applicable |
| Sponsor | Kenvue Brands LLC |
| Enrollment | 250 participants |

Key Eligibility Criteria

Inclusion (8)

- \. Adequate oral hygiene (i.e., brush teeth daily and exhibit no signs of gross oral neglect).
- \. Mean Modified Gingival Index (MGI) is greater than or equal to 2.00 at screening visit and reconfirmed at baseline visit.
- \. Bleeding upon probing (BOP) is greater than or equal to 30% at the screening visit.
- \. Diagnosis of gingivitis (confirmed by MGI and BOP) at the screening visit and reconfirmed at the baseline visit.
- \. Able to read and understand English. 7. Subject has signed the written informed consent (ICD) and indicated their agreement with the terms of the study and the study procedures listed and photograph release including the Health Insurance Portability and Accountability Act (HIPAA) disclosure prior to any study-related procedures.

... and 3 more (see full listing online)

Exclusion (28)

- Significant oral soft tissue pathology or active dental caries, based on the dentist's visual examination and at the discretion of the Investigator.
- History of significant adverse reactions, including sensitivity or suspected allergies, following use of oral hygiene products such as toothpastes, mouth rinses, and red food dye.
- Periodontitis, as determined by more than 2 sites with a probing depth greater than 4mm.
- Regular consumption of probiotics supplements within one week prior to screening. Subjects can be rescreened if they fulfill this criterion.
- Dental prophylaxis within 4 weeks prior to screening. Subjects can be rescreened if they fulfill this criterion.

... and 23 more (see full listing online)

Locations (1 total)

The ADA Forsyth Institute, Somerville, Massachusetts, United States