

# Topical Steroids to Prevent Recurrent Urinary Tract Infections in Uncircumcised Male Infants: a Pilot Study

NCT07508124

---

<b>Status</b>	RECRUITING
<b>Phase</b>	Phase 4
<b>Sponsor</b>	Baylor College of Medicine
<b>Enrollment</b>	40 participants

## Key Eligibility Criteria

---

### Inclusion (3)

- Uncircumcised male infants < 6 months old
- Both physician and laboratory confirmed diagnosis of first UTI
- Fever (temperature  $\geq 38.0$  C) or hypothermia (temperature  $< 36.0$  C) per ED/hospitalization or parental report

### Exclusion (10)

- Legal guardian not available to provide informed consent
  - Infant in Child Protective Services (CPS) custody
  - Legal guardian would like Urology referral for circumcision
  - Circumcision is medically indicated due to recurrent episodes of balanitis or ballooning of foreskin during urination
  - Past diagnosis of penile abnormalities (diagnoses of epispadias, hypospadias or congenital penile curvature (chordee))
- ... and 5 more (see full listing online)

## Locations (3 total)

---

Texas Children's Hospital Main campus, Houston, Texas, United States  
Texas Children's Hospital West Campus, Katy, Texas, United States  
Seattle Children's Hospital, Seattle, Washington, United States